




P CAIN CREAM

S CAIN CREAM

Local Anesthetic Cream



P CAIN CREAM

S CAIN CREAM

- Classified as OTC (Over the Counter) Drug
- GMP Approved



- As a local anesthetic cream, CAIN CREAM will provide the numbness to the treated area for the procedure's period.
- The main ingredient is lidocaine, which is widely used for medical needs.
- CAIN CREAM meets all the industry standards and provides effective results, which was proven by many doctors around the world.

	P CAIN CREAM	S CAIN CREAM
PACKING	500G/JAR	500G/JAR, 30g/TUBE
COMPOSITION	(25mg LIDOCAINE + 25mg PRILOCAIN)/g	105.6mg LIDOCAINE/g
PRESERVATIVE	METHYL PARAOXYBENZOATE 1.0mg/g	METHYL p-HYDROXYBENZOATE 1.0mg/g
OTHER ADDITIVES	GLYCERIN MONOSTEARATE	STEARYL ALCOHOL
	N-METHYLPYRROLIDONE	ISOPROPYL MYRISTATE
	LIQUID PARAFFIN	CARBOMER 940
	ISOPROPYL MYRISTRATE	TROLAMINE
	PURIFIED WATER	POLYSORBATE60
	CARBOMER 940	ETHANOL
SHELF-LIFE	36 MONTHS FROM MANUFACTURING DATE	36 MONTHS FROM MANUFACTURING DATE
STORAGE	ROOM TEMPERATURE	ROOM TEMPERATURE
APPEARANCE	WHITE CREAM	WHITE CREAM

P CAIN CREAM

EFFECTS Surface anesthesia of the skin during needle insertion or surgical procedures
Anesthesia of the genital mucosa (e.g. prior to remove of condyloma)
Surface anesthesia of leg ulcers prior to irrigation or topical sergery (e.g. removal of fibrin, pus or subcutaneous tissue)

DOSAGE



1. Skin
After applying this drug, cover it with sealing tape. The seal on the application site should nomally be maintained for 1-3 hours, and the effect lasts at least 1 hour after the sealing tape is removed.
2. Genital mucosa
When removing condylome, apply 10g of this drug to the treated area for 5-10 minutes, then remove this drug and start the procedure immediately.
3. Leg ulcer
After applying this drug, cover it with sealing tape. Apply 1-2g/10cm square for at least 30 minutes and do not exceed 10g for each treatment. For leg ulcers of tissues that are difficult to submit, the application time can be extended to 60 minutes, and the ulcers should be washed within 10 minutes after removal of the cream. When used up to 15 times in 1-2 months, there was no decrease in effect nor increase in local reaction.

S CAIN CREAM

EFFECTS Local anesthesia during cosmetic procedures
Reduction of sensitivity of men's sexual organ

DOSAGE Apply 0.25-0.66g (23-62mg as lidocaine) to the application site 5 to 15 minutes prior to the procedure for adults. Do not exdeed 2g (184mg as lidocaine) in 24hours.

HOW TO USE



1. Before application, wash and dry the area thoroughly.
2. CAIN CREAM should be applied directly to the skin, spread evenly over the affected area.



3. Cover with plastic wrap and keep for recommended time till complete numbness.
4. Blotting with tissue or a soft cloth before application of the product.

CAUTIONS

1. Do not apply the product around eyes as it may cause corneal damage. If the product comes into contact with eyes, the eyes should be washed with water or sodium chloride solution and should be protected until numbness returns.
2. Like other local anesthetics, it may be toxic for the ears and should not be applied to the middle ear. This product should not be used in procedures that require the penetration of middle ear
3. Do not apply to open erosions.



#522 Dongkwang BizTower, 272 Sunhwagung-ro,
Namyangju-si, Gyeonggi-do, Republic of Korea
+82-2-6225-7988

Website: www.sbodyline.com

Email: dfkbio@dfkbio.com